

Oral Presentations

- O 1 Scott S.D., Isler R.C., Murakami M., Bell J.D., Bush C.E., et al.
ORNL, Oak Ridge, Tennessee, USA
Carnevali¹ A., Murray L.E.
University of Tennessee, Knoxville, Tennessee, USA
¹Harding University, Searcy, Arkansas, USA
Neutral injection experiments on the ISX-B tokamak
- O 2 Greenwald M., Parker J., Besen M., Fiore C.L., Foord M., et al.
Plasma Fusion Center, MIT, Cambridge, Massachusetts, USA
Milora S., Schuresco D.
ORNL, Oak Ridge, Tennessee, USA
Hydrogen pellet injection into ALCATOR-C
- O 3 Burrell K.H., Doublet III Physics, Operations and Neutral Beam Groups
GA-Technologies, Inc., San Diego, Ca., USA
A regime of improved energy and particle confinement in neutral beam heated divertor discharges in DOUBLET III
- O 4 Ellis J.J., Morris A.W.
Culham Laboratory, Abingdon, Oxon, UK
Ininstability control with active saddle coils on TOSCA
- O 5 Dobrokhотов E.J., Дремин М.М., Павлов Ю.Д., Петров Д.П., Солтсев А.М., Стефановский А.М., Теребков А.Л., Шичербак А.Ф.
I.V.Kurchatov Institute of Atomic Energy,
Moscow, USSR
Bolometer measurements of the energy losses in the graphite wall tokamak
- O 6 Fußmann G., Behringer K., Bernhardi K., Haas G., Poschenrieder W., et al.
MPI für Plasmaphysik, Garching, FRG
Kissel' S., Szymanski² Z..
¹on leave from MIT, Cambridge, Massachusetts, USA
²on leave from Institute of Fundamental Technological Research, Warsaw, Poland
Impurity retainment in the divertor of ASDEX
- O 7 Meyer-ter-Vehn J., Arnold R., Metzler N., Schmalz R., Unterseer K., Witkowski S.
MPI für Quantenoptik, Garching, FRG
ICF-target research at MPQ/Garching
- O 8 Yoshida K., Miyanaga N., Kitagawa Y., Kato Y., Nishihara K., et al.
Institute of Laser Engineering, Osaka University, Osaka, Japan
Efficient and uniform compression to high density with 1.052μm GM-II laser
- O 9 Rockett P.D., Tarvin J.A., Busch Gar. E., Charatis G., Johnson R.R., et al.
KMS Fusion, Inc., Ann Arbor, Michigan, USA
The impact of quality diagnostics on ICF research at KMS-Fusion, Inc.
- O 10 Kaye S.M. Bell M., Bol K., Boyd D., Brau K., et al.
PPPL, Princeton University, Princeton, NJ, USA
High confinement discharges in PDX with neutral beam injection
- O 11 Degtyarev L.M., Poshekhanov Yu.Yu., Keldysh Institute of Applied Mathematics
Shafranov V.D.
I.V. Kurchatov Institute of Atomic Energy,
Moscow, USSR
Comparative characteristics of various plasma systems
- O 12 Chance M.S., Chen L., Furth H.P., Goldston R., Grimm R.C., et al.
PPPL, Princeton University, Princeton, NJ, USA
Dewar R.L.
Research School of Physical Sciences, A.N.U., Australia
Todd A.
Grumman Aerospace Corporation, Princeton, New Jersey, USA
Bean-shaped tokamak configurations
- O 13 Berk H.L., Rosenbluth M.N.
Institute for Fusion Studies, University of Texas at Austin, USA
Shohet J.L., Hitchon W.N.G.
University of Wisconsin, Torsatron/Stellerator Laboratory, Madison, Wisconsin, USA
Ballooning mode calculations in stellarators
- O 14 Register A., Hasselberg G., Waelbroeck F.
IPP, KFA Jülich GmbH, Jülich, FRG
On the origin of sawtooth relaxations and plasma disruptions
- O 15 Goedbloed¹ J.P., Freidberg J.P., Rohatgi R.
Plasma Fusion Center, MIT, Cambridge, Ma., USA
¹On leave from FOM-Instituut voor Plasmafysica, Rijnhuizen, Nieuwegein, The Netherlands
Influence of a limiter on the stability of external kink modes
- O 16 Faulconer D.W.
Laboratoire de Physique des Plasmas, Ecole Royale Militaire, Brussels, Belgium
Electrostatic shield fringe field interaction with plasma in rf heating
- O 17 Swain D.W., Colchin R.J., Baity F.W., Cobble J.A., Davis W.A., et al.
ORNL, Oak Ridge, Tennessee, USA
The EBT experiment
- O 18 Lyon J.F., Carreras B.A., Cooper W.A., Dory R.A., Garcia L., et al.
ORNL, Oak Ridge, Tennessee, USA
Charlton L.A., Fowler R.H., Hicks H.R., Holmes, J.A., Lynch V.E., Masden B.F.
Union Carbide Corporation Nuclear Division, Computer Sciences
ATF-1 torsatron studies
- O 19 Uo K., Iiyoshi A., Obiki T., Motojima O., Morimoto S., et al.
Plasma Physics Laboratory, Kyoto University, Kyoto, Japan
Recent results of nbi and ecr heating experiments on HELIOTRON E

VIII

- O 20 Andryukhina Eh.D., Agapov L.N., Batanov G.M., Berezhetskij M.S., Blokh M.A.. et al.
General Physics Institute, USSR Academy of Sciences, Moscow, USSR
Initial experiments on electron cyclotron plasma heating in the L-2 stellarator
- O 21 Nedospasov A.V., Tokar M.Z.
Institute of High Temperatures, USSR Academy of Sciences, Moscow, USSR
Recycling influence on plasma of a tokamak periphery
- O 22 Bernard L.C., Bhadra D.K., Helton F.J., Lao L.L.
GA Technologies, Inc., San Diego, California, USA
Todd T.N.
Culham Laboratory, Abingdon, Oxon, UK
Ideal MHD beta limits: scaling laws and comparison with DOUBLET III high-beta plasmas
- O 23 Czekaj S., Denus S., Kasperczuk A., Miklaszewski R., Paduch M., et al.
S. Kaliski Institute of Plasma Physics and Laser Microfusion, Warsaw, Poland
Investigation of connections between the plasma column evolution and the neutron emission in the plasma focus device
- O 24 Yokoyama M., Yamamoto Y., Kisoda A., Yamada Y., Kitagawa Y., Yamanaka M., Yamanaka C. et al.
Institute of Laser Engineering, Osaka University, Osaka, Japan
Dense plasma focus research in ILE, Osaka
- O 25 Herold H., Bertalot L., Jäger U., Schmidt H., Schmidt R., Shakhatre M.
Institut für Plasmaforschung, Universität Stuttgart, FRG
Plasma dynamics, neutron and ion emission of the POSEIDON plasma focus
- O 26 Brzozko¹ J.S., Conrads² H., Rager³ J.P., Robouch B.V., Steinmetz⁴ K.
Association EURATOM-ENEA, Frascati (Rome), Italy
¹University of Warsaw, Bialystok Branch, Poland
²IPP, KFA Jülich GmbH, Jülich, FRG
³DG- 12-Fusion (Meeus), CCE Brussels, Belgium
⁴MPI für Plasmaphysik, Garching, FRG
Characteristics of high energy deuterons in the Frascati 1-MJ plasma focus
- O 27 Jerzykiewicz A., Bielik M., Jankowicz Z., Kociecka K., Kocinski L., et al.
Institute for Nuclear Studies, Swierk, Poland
Borowiecki M., Czekaj S., Denus S., Skrzeczanowski W.
Institute of Plasma Physics and Laser Microfusion, Warsaw, Poland
Preliminary investigations of 360 kJ plasma-focus device
- O 28 Nardi V., Bostick¹ W.H., Powell C., Prior W.
Stevens Institute of Technology, Hoboken, N.J., USA
¹Physics Department, University of New Mexico, Albuquerque, New Mexico, USA
Ion imaging and energy spectrum from the plasma-focus ion emission
- O 29 Tuszeński M., Armstrong W.T., Barnes Cris W., Barnes D.C., Bartsch R.R., et al.
Los Alamos Natl. Lab., Los Alamos, NM., USA
Gladd N.T.
Jacor, Inc., Del Mar, California, USA
Shestakov A.I.
Lawrence Livermore Natl. Lab., Livermore, California, USA
Recent results in the Los Alamos compact torus program
- O 30 Antoni V., Buffa A., Giudicotti L., Martini S., Ortolani S., Puiatti M.E., Scarin P.
Istituto Gas Ionizzati del C.N.R., Padova, Italy
Reversed field pinch experiments at various currents in ETA-BETA II
- O 31 Watt R.G., Jacobson A.R., Massey R.S., Mondt J.P.
and the ZT-40M Team
Los Alamos Natl. Lab., Los Alamos, New Mexico, USA
Recent results from the ZT-40M reversed field pinch program
- O 32 Porkolab M., Lloyd B., Schuss J.J., Takase Y., Texter S., et al.
Plasma Fusion Center , MIT, Cambridge, Ma., USA
Lower hybrid current drive and heating experiments at the 1MW rf power level on ALCATOR-C
- O 33 Nagayama Y., Tsuji S., Miyamoto K., Department of Physics, University of Tokyo, Tokyo, Japan
Kawahata K., Noda N., Tanahashi S., Fujita J., Institute of Plasma Physics, Nagoya University, Nagoya, Japan
A study of tokamak disruptions in JIPP T-II using soft X-ray imaging

- E 44 Wooding E.R., Brown D.A., Marchington P.S.,
Sayed N., Sudera Y.
Royal Holloway College, University of London, UK
Density profiles and fluctuations in CO₂ laser produced plasmas
- E 45 Murusidze I.G., Tsintsadze N.L. Tskhakaya D.D.,
Institute of Physics, Academy of Sciences of the Georgian SSR, Tbilisi, USSR
Radiation of Alfvén waves by the accelerating Langmuir soliton

Diagnostics

Part II, p. 51 - 100

Laser-Plasma Interaction, Intertial Confinement
Part II, p. 1 - 50

- E 37 Atzeni S., Caruso A.
CRE, Frascati, Italy
Ignition criterion for dense, isobaric D-T plasmas and consequences on ICF target design
- E 38 Heikkilä J.A., Karttunen S.J., Salomaa¹ R.R.E.,
Technical Research Centre of Finland,
Nuclear Engineering Laboratory, Helsinki, Finland
¹Helsinki University of Technology, Espoo, Finland
Nonlinear ion waves in absolute Brillouin scattering
- E 39 Yoshida K., Kato Y., Nakatsuka M.,
Jitsuno T., Nowada S. et al.
Institute of Laser Engineering, Osaka University, Suita, Osaka, Japan,
GEKKO XII glass laser system for laser fusion research
- E 40 Hora H., Lalosius P., Ghatak A.K., Jones¹ D.A.
Telapaga-Rosenberger S.
Department of Theoretical Physics, University of New South Wales, Kensington, Australia
¹Materials Research Labs, Ascot Vale, Victoria
High electrostatic fields and oscillations in expanding inhomogeneous plasmas without external fields or with laser irradiation
- E 41 Sternlieb A.
Soreq Nuclear Research Centre, Yavne, Israel
Scaling laws for the resistive longitudinal instability in finite charged beams
- E 42 Nicolas J.A., Sanmartin J.R.
ETSI Aeronáuticos, Universidad Politécnica de Madrid, Spain
Inverse Bremsstrahlung effects on the coronae of laser-irradiated pellets and slabs
- E 43 Barrero A., Rosety L., Sanz¹ J.
ETSI Industriales, Universidad de Sevilla, Spain
ETSI Aeronáuticos, Universidad de Politécnica de Madrid, Spain
Spherical expansion plasmas in ion beam impinged pellets
- E 14 Mori I.
Technical College of Tokushima University, Tokushima, Japan
Ion temperature measurement with neutral beams
- E 15 Alladio F., Crisanti F.
CRE, Frascati, Italy
Equilibrium magnetic measurements by toroidal multipolar method for FTU tokamak
- E 16 Lehner Th.
Laboratoire de Physique des Milieux Ionisés, Ecole Polytechnique, Palaiseau, France
Ion Bernstein modes fluctuations as a measure of the magnetic field orientation
- E 17 von Hellermann M., Gükler H.J., Hasse O.R., FB Physik, Universität Essen, FRG
Observation of plasma turbulence by coherent far forward scattering using a multi channel detector array at 10 microns
- E 18 Airolidi A.C., Oreifice A., Ramponi G.
Istituto di Fisica del Plasma, CNR Milano, Italy
Ece simulation for a toroidal plasma and its applications to JET
- E 19 Hallcock G.A., Hickok R.L., Jennings W.C., Connor K.A., Forester J.C. et al.
Rensselaer Polytechnic Institute, Troy, NY., USA
Haste G.R., Wootton A.J.
ORNL, Oak Ridge, Tennessee, USA
Space potential measurements on EBT-S and ISX-B
- E 20 Tonetti G.
CRPP-EPFL, Lausanne, Switzerland
Measurement of the plasma magnetic field with an ion beam probe
- E 21 Woskoboinikow P., Mulligan W.J., Machuzak J., Cohn D.R., Temkin R.J. et al.
Plasma Fusion Center, MIT, Cambridge, USA
385μm D₂O Laser collective Thomson scattering ion temperature diagnostic
- E 22 Gimzewski J.K., Veprek S.
Institute of Inorganic Chemistry, University of Zürich, Zürich, Switzerland
A new method for the determination of impurity ion energies and their implantation depths in collection probes exposed to the scrape-off layer

- E 23 Lisitano G., Baretich¹ F., Gammino¹ G., Bazzo I., Cassamagnagh¹ C. et al.
MPI für Plasmaphysik, Garching, FRG
¹Cise, Department of Acoustical Diagnostics, Milano, Italy
Millimeter wave Schlieren diagnostics for fusion plasma
- E 24 W VII-A Team: Tutter M., Cattanei G., Dorst D., Elsner A. Erckmann V. et al.
NI-Group: Freudenberger K., Lister G.G., Ott W., Penningfeld F.-P., Speth E.
MPI für Plasmaphysik, Garching, FRG
Effect of suprathermal electrons on Ece-spectra in W VII-A
- E 25 Filippov V.N., Komissarova I.I., Ostrovskaya G.V., Shedova E.N.
A.F. Ioffe Physico-Technical Institute, Leningrad, Academy of Sciences, USSR
The prospects to enhance the sensitivity of holographic interferometry plasma diagnostics in IR-region
- Magnetic Confinement Theory**
Part II, p. 101 - 288
- B 41 Ming-Lun Xue
Institute of Mechanics, Chinese Academy of Sciences, Beijing, People's Republic of China
Advanced concept of adiabatic compression of tokamak and self-compression of compact torus
- B 42 Khait V.D., Kolesnikov V.K.
Institute of High Temperatures, USSR Academy of Sciences, Moscow, USSR
Solution of the tokamak plasma equilibrium at high betas by the variational method
- B 43 Casci F., Kaleck¹ A., Zehrfeld H.P.
MPI für Plasmaphysik, Garching, FRG
¹IPP, KFA Jülich GmbH, Jülich, FRG
Quasistatic evolution of ideal MHD equilibria in INTOR and ASDEX-UG
- B 44 Bernardin M.P., Tataronis J.A.
University of Wisconsin, Madison, Wisconsin, USA
Force-free MHD equilibria in non-axisymmetric toroidal geometry
- B 45 Kovriznykh L.M., Shchepetov S.V.
Institute of General Physics, Moscow, USSR
Condition for the stability of dissipative modes in a stellarator
- B 46 Rem J., Galvao¹ R.M.O., Goedbloed J.P., Sakanaka² P.H.
FOM-Instituut voor Plasmafysica, Rijnhuizen, Nieuwegein, The Netherlands
¹Centro Tecnico Aerospacial, São Paulo, Brazil
²Universidade Estadual de Campinas, São Paulo, Brazil
Ideal MHD stability of a tokamak in the high-beta tokamak ordering
- B 47 Nührenberg J., Großmann R., Herrnegger F.
MPI für Plasmaphysik, Garching, FRG
Evaluation of Mercier's criterion in 3D HELIAC equilibria
- B 48 Pogutse O.P., Yurchenko E.I.
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
Critical plasma pressure provided by the ballooning modes of helical instability in tokamak
- B 49 Pogutse O.P., Chudin N.V.
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
Radial structure of resistive ballooning modes
- B 50 Rosenau Ph.
Faculty of Mechanical Engineering, Israel Institute of Technology, Haifa, Israel
On tearing modes in a resistive medium
- B 51 Saramito B., Maschke E.K.
Département de Recherches sur la Fusion Contrôlée, CEN, Fontenay-aux-Roses, France
Bifurcation theory for tearing modes
- B 52 Dnestrovskij¹ Yu.N., Kosomarov D.P., Popov A.M., Shagirov E.A.
M.V. Lomonosov State University, Moscow, USSR
¹I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
Non-linear evolution of helical modes in a tokamak with low q
- B 53 Samain A.
Département de Recherches sur la Fusion Contrôlée, CEN, Fontenay-aux-Roses, France
Diamagnetic destabilization of magnetic islands in the non-linear regime
- B 54 Nedospasov A.V.
Institute of High Temperatures, USSR Academy of Sciences, Moscow, USSR
Current-convective turbulence and anomalous diffusion in tokamaks
- C 37 Evangelidis E.A.
Plasma Physics Division, NUCOR-Pelindaba Pretoria, South Africa
On the solution of the equation of plasma equilibrium in a toroidal configuration
- C 38 Mikhailov M.I.
"Energiya" Scientific Manufacturing Association Shafrarov V.D.
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
Variational principle in flux variables for the toroidal plasma equilibrium
- C 39 Wahlberg C., Arora¹ A.K., Ring. R.
Institute of Technology, Uppsala, Sweden
¹Department of Physics, University of Rajasthan, Jaipur, India
Z-Pinch Equilibria in an External Magnetic Multipole Field
- C 40 Grauer R., Rebhan, E.
Institut für Theoretische Physik II, Universität Düsseldorf, Düsseldorf, FRG
Semianalytical calculation of equilibrium and axisymmetric stability of finite aspect ratio tokamaks with elliptic cross-section
- C 41 Kratzsch, K.A., Rebhan E.
Institut für Theoretische Physik II, Universität Düsseldorf, Düsseldorf, FRG
Calculation of growth rates from the marginal mode

XVII

- C 42 Rewoldt G., Johnson¹ J.L., Anania² G., Chance M.S., Manickam J. et al.
PPPL Princeton University, New Jersey, USA
'On loan from Westinghouse Research & Development Center
²At Booz Allen and Hamilton, Management Consultants
Application of the stellarator expansion to equilibrium and stability studies of three-dimensional toroidal configurations
- C 43 Kammarsh T., Nguyen K., University of Michigan, Ann Arbor, Michigan, USA
Beta limit for ballooning instabilities in EBT
- C 44 Cheng C.Z., Chen L., Chance M.S., Rewoldt G., Tang W.M.
PPPL Princeton University, Princeton, New Jersey, USA
Shear Alfvén waves and ballooning modes in tokamaks
- C 45 Sundaram A.K., Sen A., Kaw P.K.
Physical Research Laboratory, Ahmedabad, India
Resistive ballooning modes in a high-temperature plasma
- C 46 Storer R.G.
The Flinders University of South Australia, Bedford Park, South Australia
Resistive magnetohydrodynamic spectrum for toroidal plasmas
- C 47 Tagger M., Frey M.
Departement de Recherches sur las Fusion Contrôlée, CEN, Fontenay-aux-Roses, France
Pellet R.
Ecole Polytechnique, Palaiseau, France
Quasi-linear evolution of tearing modes
- C 48 Gribkov V.M., Kadomtsev B.B., Pogutse O.P.
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
Plasma pressure drops produced by the ballooning modes
- C 49 Lee J.K., Chu M.S., Helton F.J.
GA Technologies, Inc., San Diego, California, USA
Park W.
PPPL, Princeton University, Princeton, New Jersey, USA
Magnetic fluctuations in high- β tokamak plasmas
- D 45 Klieb L., Schep T.J., Venema M.
FOM-Instituut voor Plasmafysica, Rijnhuizen, Nieuwegein, The Netherlands
Pegoraro F.
Scuola Normale Superiore, Pisa, Italy
Collisional and kinetic effects on electromagnetic modes in toroidal geometry
- D 46 Schep T.J., Venema M.
FOM-Instituut voor Plasmafysica, Rijnhuizen, Nieuwegein, The Netherlands
Drift modes in toroidal geometry
- D 47 Pegoraro F., Coppi¹ B., Ramos J.
Scuola Normale Superiore, Pisa, Italy
¹MIT, Cambridge, Massachusetts, USA
Electromagnetic ion cyclotron modes in a plasma with polarized nuclear spins
- D 48 Hasselberg G., Register A.
IPP, KFA Jülich GmbH, Jülich, FRG
Drift wave spectra, energy fluxes and impurity transport in tokamak plasmas
- D 49 Andersson P.
Institute for Electromagnetic Field Theory, Chalmers University of Technology, Göteborg, Sweden
The effects of temperature gradients on the curvature driven trapped electron mode
- D 50 Hamnén H., Bondeson A., Anderson D., Lisak M., Core¹ W.G.F.
Institute for Electromagnetic Field Theory, Chalmers University of Technology, Göteborg, Sweden
¹JET Joint Undertaking, Abingdon, Oxon, UK
Ripple-induced radial diffusion of fast ions in tokamaks
- D 51 Weiland J.
Institute for Electromagnetic Field Theory, Chalmers University of Technology, Göteborg, Sweden
A nonlinear gyrokinetic equation obtained by integration along unperturbed orbits
- E 46 Nocentini A., De Meio L., Istituto di Meccanica dell' Università Trieste, Italy
A new pseudo-classical transport theory for tokamak plasmas
- E 47 Thyagaraja A., Haas F.A.
Culham Laboratory, Abingdon, Oxon, UK
A novel test-particle approach to anomalous transport in tokamaks
- E 48 Gurevich A.V., Dimant Ya.S., Dnestrovskij¹ Yu.M., Kostomarov², D.P., Smirnov² A.P.
Lebedev Institute of Physics, Moscow, USSR
¹I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
²M.V. Lomonosov State University, Moscow, USSR
Simulation of convective electron transport in T-10 tokamak
- E 49 Dnestrovskij Yu.N., Neudatchin S.V. Pereverzev G.V., Polovoj A.R.
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
Anomalous pinching and particle balance in tokamak
- E 50 Yushmanov P.N.
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
Generalized ripple-banana transport in tokamak
- E 51 Yushmanov P.N.
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
High energy ion confinement in tokamaks with local divertor
- E 52 Avinash K.
Physical Research Laboratory, Navrangpura, Ahmedabad, India
Is motion of charged particle in mirror of true Markovian nature?
- E 53 Riedler J.M., Heindler M., Miley¹ G.H.
Institut für Theoretische Physik, TU Graz, Austria
¹Nuclear Engineering Laboratory, Urbana, Illinois, USA
Coulomb scattering between Maxwellian plasma and monoenergetic beam

Transport codes

Part II, p. 289 - .366

- A 12 Rau F., Chodura R., Kisslinger J., Lotz W., Wobig H.
MPI für Plasmaphysik, Garching, FRG
Some effects of finite plasma pressure in the advanced stellarator WENDELSTEIN VII-AS
- A 13 Nicolai A., Reiter D.
IPP, KFA Jülich GmbH, Jülich, FRG
Two-dimensional computer modelling of JET as a specific example of D-shaped tokamak plasmas
- A 14 Becker G., Behringer K., Bernhardi K., Ebernhagen A., Fußmann G., et al.
MPI für Plasmaphysik, Garching, FRG
Kissel¹ S., Szymanski² Z..
¹On leave from MIT, Cambridge, Massachusetts, USA
²On leave from Institute of Fundamental Technological Research, Warsaw, Poland
Transport modelling of neutral beam heated I and h discharges in ASDEX
- A 15 Watkins M.L.
JET Joint Undertaking, Abingdon, Oxon, UK
Abels van Maanen A.E.P.M.
On attachment from FOM-Instituut voor Plasmafysica, Rijnhuizen, Nieuwegein, The Netherlands
Tokamak transport calculations for JET with different limiter materials
- A 16 Sidorov V.P., Minenkov V.P., Komoshvili K.G., Vekua Institute of Physics and Technology, Sukhumi, USSR
Favorsky A.P., Dmitrieva M.V., Tishkin V.F.
Keldysh Institute of Applied Mathematics, Moscow, USSR
Numerical calculations of plasma parameters for a steady-state Alfvén-wave tokamak
- A 17 Pavlichenko D.S., Besedin N.T., Pyatov V.N., Slavyi A.S., Shvets O.M.
Institute of Physics and Technology, Ukrainian Academy of Sciences, Kharkov, USSR
Neoclassical equilibrium of plasma during ion heating in stellarator
- A 18 Core W.G.F.
Jet Joint Undertaking, Abingdon, Oxon, UK
A 2-d model for the calculation of neutral transport in tokamaks with non-circular magnetic surfaces
- A 19 Nicolai A., Wolf G.H., IPP, KFA Jülich GmbH, Jülich, FRG
Modelling of tokamak discharges based on transport processes along oblique magnetic field lines
- A 20 Lister G.G., Penningsfeld F.P., Ott W., Speth E.
MPI für Plasmaphysik, Garching, FRG
Neutral beam injection studies for W VII AS using the 3-d computer code FAFNER
- A 21 Watkins M.L., Düchs D.F., Noll P.
JET Joint Undertaking, Abingdon, Oxon, UK
Is additional edge transport beneficial?
- B 15 Sano F., Obiki T., Hanatani K., Zushi H., Sudo S. et al.
Plasma Physics Laboratory, Kyoto University, Gokasho, Uji, Kyoto, Japan
Ion and electron energy balance analysis of neutral-beam-heated Heliotron E plasmas
- B 16 Martinez-Val J.M., Velarde G., Aragonés J.M., Arroyo R., Gago J.A. et al.
Departamento de Energía Nuclear E.T.S.I. Industriales Universidad Politécnica de Madrid, Spain
Analysis of charged-particle energy deposition for ICF calculations with the NORCLIA Code
- B 17 Minardi E.
Istituto di Fisica del Plasma, Associazione EUR-CNR, Milano, Italy
Plasma circuit instability and current disruption
- B 18 Tendler M., Neuhauser¹ J., Wunderlich¹ R.
Royal Institute of Technology, Stockholm, Sweden
¹MPi für Plasmaphysik, Garching, FRG
Numerical simulations of the oxygen impurity transport at the plasma periphery in a tokamak
- B 19 Tendler M.
Royal Institute of Technology, Stockholm, Sweden
EXTRAP in the quasi-stationary regime
- B 20 Lindemann¹ E.L., Rickard G., Haines M.G.
Blackett Laboratory, Imperial College, London, UK
¹On leave from Los Alamos National Laboratory, New Mexico, USA
An implicit "quasineutral" plasma simulation method which includes sheath and vacuum regions
- B 21 Hitchon W.N.G., Myrick H.E.
Torsatron / Stellarator Lab. University of Wisconsin, Madison, Wisconsin, USA
A model for the magnetic field strength along a stellarator field line and its application to particle orbit calculations
- B 22 Turner M.F.
Culham Laboratory, Abingdon, Oxon, UK
Wesson J.A.
JET Joint Undertaking, Abingdon, Oxon, UK
JET simulation
- B 23 Sykes A., Turner M.F., Patel S.
Culham Laboratory, Abingdon, Oxon, UK
 β -limits in tokamaks due to high-n ballooning modes

- Plasma-Wall Interaction, Divertors**
Part II, p. 367 - 504
- A 32 Krüger H., Rödiger K., Schulz V., Stampa A.
FB Physik, Universität Essen, FRG
Experiments on plasma-induced arcs
- A 33 Ertl K., Behrisch R., ASDEX Team and PWW Team
MPI für Plasmaphysik, Garching, FRG
Jüttner, B.
Zentralinstitut für Elektronenphysik, Akademie der
Wissenschaften der DDR, Berlin, GDR
**Investigation of electric arcs in the divertor of
the tokamak experiment ASDEX**
- A 34 Heifetz¹ D., Lackner K., Rauh K.G., Schneider W.
MPI für Plasmaphysik, Garching, FRG
¹PPPL, Princeton, New Jersey, USA
**Simulation of ASDEX divertor action using a realistic
neutral gas model**
- A 35 Wienhold P., Waelbroeck F., Winter J., Rota E.,
Banno T.
IPP, KFA Jülich GmbH, Jülich FRG
**In situ measurement of the hydrogen recycling
constant of the TEXTOR liner**
- A 36 Iglyanov Yu.L., Kukushkin A.S., Pigarov A.Yu.,
Pistunovich V.I.,
I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR
**Two-dimensional calculations of near-wall plasma
dynamics for a tokamak with a poloidal divertor**
- A 37 Petrov V.G.
Institute of High Temperatures, USSR Academy
of Sciences, Moscow, USSR
**The characteristics of stationary plasma flow in the
tokamak scrape-off layer**
- A 38 Zykova N.M., Ikonnikov A.O., Kabanov V.N.,
Nedospasov A.V.
Institute of High Temperatures, USSR Academy
of Sciences, Moscow, USSR
Arcing study in disruptive tokamak discharges
- A 39 Fidel'man G.N., Ivanov R.S., Stotsky G.J.,
Institute of High Temperatures, USSR Academy of
Sciences, Moscow, USSR
**On longitudinal transport of charged particles
to tokamak limiter**
- A 40 Samm U.
IPP, KFA Jülich GmbH, Jülich, FRG
**Results from infrared-thermography on limiters
in TEXTOR**
- A 41 Pospieszczyk A., Bogen P., Samm U.
IPP, KFA Jülich GmbH, Jülich, FRG
**Neutral particle fluxes emitted from the TEXTOR
limiter**
- A 42 Betz G., Husinsky W., Viehböck, F.
Institut für Allgemeine Physik, TU Wien, Wien, Austria
**Influence of oxygen coverage on the sputtering of
chromium**
- B 32 Sugai H.
Department of Electrical Engineering, Nagoya University,
Nagoya, Japan
Radio-frequency divertor for impurity pump-out
- B 33 Harbour P.J., Morgan J.G.
Culham Laboratory, Abingdon, Oxon, UK
**The transport of impurity ions in a scrape-off
plasma**
- B 34 Braams B.J.
FOM-Instituut voor Plasmafysica, Rijnhuizen,
Nieuwegein, The Netherlands
**Numerical studies of the two-dimensional
scrape-off plasma**
- B 35 Fielding S.F., Mc Cracken G.M.
Culham Laboratory, Abingdon, Oxon, UK
**A comparison of impurity levels in JET with nickel,
carbon and beryllium limiters**
- B 36 Harrison M.F.A., Hotston E.S.
Culham Laboratory, Abingdon, Oxon, UK
**Exhaust of helium and impurity release in INTOR with
a divertor or a pumped limiter**
- B 37 Gruber O., Lackner K.
MPI für Plasmaphysik, Garching, FRG
**Pooidal field requirements for divertor and
pump limiter design**
- B 38 Daybelge U.
Marmara Research Institute, Department of Applied
Mathematics, Gebze, Kocaeli-Turkey
**Asymmetry of plasma scrape-off in limiter and
divertor tokamaks**
- B 39 Clark W.H.M., Fielding S.J., Goodall D.H.J.,
Hugill J., Johnson P.C. et al.
Culham Laboratory, Abingdon, Oxon, UK
Energy and power balances in DITE tokamak
- B 40 Fielding S.J., Goodall D.H.J., Hugill J., Johnson P.C.
Culham Laboratory, Abingdon, Oxon, UK
**Characteristics of the particle and energy exhaust
into the DITE bundle divertor**
- C 25 Shimada M., Sengoku S., Kobayashi T., Miya N.,
Kasai M. and the JAERI Team
JAERI, Tokai, Japan
Kahn C.L., Burrell K.H. and the GA Doublet III Groups
GA Technologies, Inc., San Diego, California, USA
**Experimental observation of a dense and cold
divertor plasma in D-III beam-heated divertor
discharges and its numerical simulation**
- C 26 Gimzewski¹ J.K., Veprek¹ S., Hofmann F.,
Lister J.B., Pochelon A., Groner² P.
CRPP-EPFL, Lausanne, Switzerland
¹Institute of Inorganic Chemistry, University of
Zürich, Switzerland
²Federal Institute for Reactor Research Würenlingen,
Switzerland
**Impurity fluxes in the scrape-off layer of the TCA
tokamak: effects of rf heating**
- C 27 Behringer K., Fußmann G., Poschenrieder W.,
Taglauer E., Bernhardi K., et al.
MPI für Plasmaphysik, Garching, FRG
Kissel¹ S., Szymanski² Z.,
¹On leave from MIT, Cambridge, Massachusetts, USA
²On leave from Institute of Fundamental
Technological Research, Warsaw, Poland
**Production and transport of impurities in the
ASDEX tokamak plasma**
- C 28 Joye B., Behn R., Hofmann F., Hollenstein Ch.,
Lister J.B. et al.
CRPP-EPFL, Lausanne, Switzerland
¹University of Fribourg, Switzerland
Peacock N.J., Stamp M.F.,
Culham Laboratory, Abingdon, Oxon, UK
**A comparison between steel, carbon and TiC-coated
limiters in the TCA tokamak**

- | | | | |
|---|---|------|--|
| C 29 | Neuhäuser J., Schneider W., Wunderlich R. MPI für Plasmaphysik, Garching, FRG Impurity flow in the tokamak scrape-off layer | D 2 | Dabiri A.E., Glancy J.E., Science Application, Inc., La Jolla, California, USA d-d compact reversed field pinch reactor technology assessment |
| C 30 | Chodura R. MPI für Plasmaphysik, Garching, FRG Plasma transport to the wall through the electrostatic sheath | D 3 | Gill R.D., JET Joint Undertaking, Abingdon Oxon, UK Empirical determination of the parameters of ignition and reactor tokamaks |
| C 31 | Winter J., Waelbroeck F., Wienhold P., Esser H.G., Könen L. et al. IPP, KFA Jülich GmbH, Jülich, FRG 'ORNL, Oak Ridge, Tennessee, USA Surface conditioning of the TEXTOR liner and limiters by plasmachemical carbon deposition | D 4 | Sternlieb A., Soreq Nuclear Research Centre, Yavne, Israel The Mirrotron – a compact disposable tandem mirror fusion reactor |
| C 32 | Artemenkov L.J., Gurov A.A., Kován J.A., Maltsev S.G., Melikhov P.J. et al. I.V. Kurchatov Institute of Atomic Energy, Moscow, USSR Experiments in TO-2-tokamak with toroidal divertor | D 5 | Lodder, J., FOM-Instituut voor Plasmafysica, Rijnhuizen, Nieuwegein, The Netherlands Nuclear spin depolarization by classical mixing in fusion reactor plasmas |
| C 33 | Alexander K.F., Grote H., Hildebrandt D., Laux M., Kota'í E. et al. Zentralinstitut für Elektronenphysik, Akademie der Wissenschaften der DDR, Berlin, GDR 'Central Research Institute for Physics, Hungarian Academy of Sciences, Budapest, Hungary Deuterium and impurity fluxes in the limiter shadow of T-10: indication for time-dependent local sources | D 6 | Kernbichler W., Niederl K., Heindler M., Technische Universität Graz, Austria Physics and engineering constraints on the feasibility of advanced fuel fusion reactors |
| C 34 | Grosman A., TFR Group Département de Recherches sur la Fusion Contrôlée, CEN, Fontenay-aux-Roses, France Plasma edge measurements during additional heating in the TFR tokamak | D 7 | Carretta, U., CNR-Politecnico di Milano, Italy Farina D., Lontano M., Minardi E., Pozzoli R., Istituto di Fisica del Plasma, CNR-Milano, Italy Burn optimization in a thermonuclear reactor |
| C 35 | Kever H. IPP, KFA Jülich GmbH, Jülich, FRG Scaling of plasma scrape-off layers in tokamaks | D 8 | Wobig H., Broßmann U., Gorenflo H., Harmeyer E., Kisslinger J. et al. MPI für Plasmaphysik, Garching, FRG Some aspects of an advanced stellarator reactor |
| C 36 | Johnson P.C., Mudford B.S., Cordey ¹ J.G. Culham Laboratory, Abingdon, Oxon, UK 'JET Joint Undertaking Neutral beam injection modelling for DITE with the MKII bundle divertor | D 9 | Bobbio S., Coccoresi V., Fabricatore G., Martone R., Rubinacci G., Istituto di Eletrotecnica, University of Naples, Italy 'Dipartimento Elettrico, University of Calabria, Italy Passive stabilization in INTOR reactor |
| Reactor Aspects Part II, p. 505 - 546 | | | |
| D 1 | Brunelli B., Zampaglione V. CRE, Frascati, Italy Avanzini P.G., Raia G., Rosatelli F. NIRA - C.P. Genoa, Italy HTMR- a hybrid toroidal magnet tokamak reactor study | D 10 | 'Brzosko J., Ingrosso L., Robouch B.V., EURATOM-ENEA, Frascati, Rome, Italy 'On leave from Warsaw University, Biocyklik Branch, Poland Depression of the nuclear properties of fusion reactor breeding blankets by nonhomogeneities |