



# CONTENTS

Introduction . . . . .	vii
Table I. The Normal Probability Integral $\Phi_0(x) = (1/\sqrt{2\pi}) \int_0^x e^{-\frac{1}{2}t^2} dt$ and its Derivative $\Phi'_0(x) = \varphi(x) = (1/\sqrt{2\pi}) e^{-\frac{1}{2}x^2}$ , $x = 0.000 (0.001) 2.500 (0.002) 3.400 (0.005) 4.00 (0.01) 4.50$ . . . . .	1
Table II. The Normal Probability Integral $\Phi_0(x) = (1/\sqrt{2\pi}) \int_0^x e^{-\frac{1}{2}t^2} dt$ and its Derivative $\Phi'(x) = \varphi(x) = (1/\sqrt{2\pi}) e^{-\frac{1}{2}x^2}$ , $x = 4.50 (0.01) 6.00$ . . . . .	35
Table III. The Function $-\log_{10} Q(x) = -\log_{10} [\frac{1}{2} - \Phi_0(x)]$ , $x = 5 (1) 50 (10) 100 (50) 500$ . . . . .	37
Table IV. The Tetrachoric Functions $\tau_s(x) = \frac{(-1)^{s-1}}{\sqrt{s!}} \frac{d^{s-1}}{dx^{s-1}} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2},$ being Normalized Derivatives of $\varphi(x) = (1/\sqrt{2\pi}) e^{-\frac{1}{2}x^2}$ , $s = 2, 3, \dots, 21$	39
Table V. The coefficients $p_s$ and $q_s$ in the Recurrence Relation $\tau_s(x) = xp_s \tau_{s-1}(x) - q_s \tau_{s-2}(x),$ $s = 2 (1) 21$ . . . . .	121
Table VI. The coefficients $\lambda_s = \sqrt{s!}$ , $s = 1 (1) 25$ , . . . . .	123
Table of Coefficients for Bessel Quadratic Interpolation . . . . .	125
List of Volumes in the Mathematical Tables Series . . . . .	127