

CONTENTS

	<i>Preface</i>	ix
1	Ancient History of Free-Electron Devices	1
	<i>E. T. Jaynes</i>	
	We recall some early thinking about the production of light from high-energy electrons that started with problems in the design of electron accelerators.	
2	Nonlinear Theory of Free-Electron Devices	41
	<i>Frederic A. Hopf, Pierre Meystre, Gerald T. Moore, and Marlan O. Scully</i>	
	Gain, saturation, recoil and velocity spread in free-electron devices are derived classically.	
3	The Free-Electron Laser as a Travelling Wave Amplifier	115
	<i>Norman M. Kroll</i>	
	Classical travelling wave tube analysis is extended to include the regime of the Stanford free-electron laser.	
4	One-Body Analysis of Free-Electron Lasers	157
	<i>William B. Colson</i>	
	In a free-electron laser electrons travel through a static, spatially periodic transverse magnetic field and oscillate to amplify radiation present in a resonant cavity. The basic gain mechanism and electron-dynamics are described by relating the phase space paths for electrons to those of a simple pendulum.	
5	Collective and Single Electron Interactions and Cerenkov-Smith-Purcell Free-Electron Lasers	197
	<i>Abraham Gover and Amnon Yariv</i>	
	A simple general model delineates the different operational regimes. Microwave tube theory is compared with the laser physics description of free-electron devices, Smith-Purcell and Cerenkov radiation and travelling wave amplifiers.	

6	Stimulated Backscattering from Relativistic Unmagnetized Electron Beams	241
	<i>Phillip A. Sprangle</i>	
	An analysis of stimulated scattering of a high frequency incident pump wave from an unmagnetized relativistic electron beam.	
7	Gyrotrons and Dopplertrons--Electromagnetic Wave Generators Based on Coherent Radiation from Magneto- Resonant, Relativistic, Electron Beams	273
	<i>Victor L. Granatstein</i>	
	The cyclotron maser and beam-wave scattering have been used to generate intense radiation of millimeter and submillimeter wavelengths.	
8	Stimulated Electromagnetic Shock Radiation (SESR)	301
	<i>Stanley Schneider and Richard Spitzer</i>	
	A new effect is predicted capable of producing intense, narrowband, tunable electromagnetic radiation by the interaction in a polarizable medium of coherent electromagnetic waves with charged particles.	
9	Stimulated Cerenkov Radiation	357
	<i>John E. Walsh</i>	
	The recent availability of intense relativistic electron beams has made possible substantial beam power, in the wavelength range, via Cerenkov interaction. Particular attention is given to stimulated emission from relativistic electron beams propagating in a variety of dielectric resonators.	
10	Generation of X-Radiation Via Electrons in a Crystal: Nonlinear Rise and Line Narrowing	381
	<i>K. Das Gupta</i>	
	A nonlinear rise in intensity and narrowing of copper $K\alpha_1$ and $K\alpha_2$ lines were observed when copper crystals were irradiated by a micro-focus high density electron beam.	
	Index	399