

Contents

Part I Astrophysical Background and Light Scattering Theory

1	Prologue	3
2	Solid Particles in Space	5
2.1	Evidence for interstellar dust	5
2.2	Observed data on optical properties of dust	6
2.2.1	Interstellar extinction	6
2.2.2	Mass density and solid particle character of absorber	10
2.2.3	Equation of transfer	12
2.2.4	Interstellar polarization	12
2.2.5	Diffuse galactic light	15
2.2.6	Interstellar dust theories	17
3	Light Scattering by Spherical Particles	25
3.1	Mie formulae	25
3.2	Computation procedure for evaluating Mie formulae	26
3.3	Asymptotic formulae for homogeneous spheres	29
3.4	Composite spheres	29
3.5	Asymptotic formulae for composite spheres	31
4	Light Scattering by Cylinders and Ellipsoids	33
4.1	Rigorous formulae for cylinders	33
4.2	Approximate formulae for infinite cylinders	34
4.3	Rayleigh scattering by ellipsoids	35

Part II Numerical Results

5	Light Scattering Functions for Homogeneous Spheres	41
	Tables T1.1–T1.169: Numerical values of $Q_{ext}(x)$, $Q_{sca}(x)$, $Q_{pr}(x)$, $Q_{bk}(x)$ and $g(x)$ for spheres characterized by a refractive index $m=n-ik$ spanning various ranges.	42
	Figs. A1–A64: $Q_{ext}(x)$, $Q_{sca}(x)$ for representative cases including examples of $g(x)$, $Q_{bk}(x)$.	211
	Figs. B1–B20: Phase functions $I_1 = S_1 ^2$, $I_2 = S_2 ^2$ as functions of θ for representative values of m and for $x=2\pi a/\lambda=1.2, 3.0, 6.0, 9.0, 12.0$.	233
6	Light Scattering Functions for Infinite Cylinders	243
	Tables T2.1–T2.104: Numerical values of $Q_{eE}(x)$, $Q_{eH}(x)$, $Q_{sE}(x)$, $Q_{sH}(x)$ for infinite cylinders at normal incidence characterized by a complex refractive index $m=n-ik$ spanning various ranges.	244
	Figs. C1–C22: Q_{eE} , Q_{eH} , Q_{sE} , Q_{sH} and $(Q_{eE} - Q_{eH})$ for infinite cylinders as functions of x for selected cases of normal incidence ($\theta=90^\circ$).	348
	Tables T2.105–T2.164: Numerical values of $Q_{eE}(x)$, $Q_{eH}(x)$, $Q_{sE}(x)$, $Q_{sH}(x)$ for infinite cylinders at oblique incidence θ .	356
	Figs. D1–D33: Q_{eE} , Q_{eH} and $(Q_{eE} - Q_{eH})$ for infinite cylinders as functions of x for selected cases of oblique incidence.	416

7	Light Scattering Functions for Spheres of Ice, Iron and Graphite	429
	Fig. E1: Complex refractive indices of ice, iron and graphite used in the present calculations.	430
	Tables T3.1–T3.20: Numerical values of Q_{ext} , Q_{sca} , Q_{bk} , g , Q_{pr} as functions of wavenumber ($x = \lambda^{-1}$ (μ^{-1})) for ice spheres of radii $a = 0.05$ (0.05) 1.0 μ , $\lambda^{-1} = 0.6$ (0.1) 10.0 μ^{-1} .	431
	Tables T3.21–T3.45: Numerical values of Q_{ext} , Q_{sca} , Q_{bk} , g , Q_{pr} as functions of wavenumber ($x = \lambda^{-1}$ (μ^{-1})) for iron spheres of radii $a = 0.01$ (0.01) 0.7 μ , $\lambda^{-1} = 0.3$ (0.1) 3.8 μ^{-1} .	451
	Tables T3.46–T3.70: Numerical values of Q_{ext} , Q_{sca} , Q_{bk} , g , Q_{pr} as functions of wavenumber ($x = \lambda^{-1}$ (μ^{-1})) for graphite spheres of radii $a = 0.01$ (0.01) 0.7 μ , $\lambda^{-1} = 0.4$ (0.1) 8.4 μ^{-1} .	464
	Figs. E2–E11: $Q_{ext}(\lambda^{-1})$, $Q_{sca}(\lambda^{-1})$ for ice spheres.	489
	Figs. E12–E16: $Q_{ext}(\lambda^{-1})$, $Q_{sca}(\lambda^{-1})$ for iron spheres.	492
	Figs. E17–26: $Q_{ext}(\lambda^{-1})$, $Q_{sca}(\lambda^{-1})$ for graphite spheres.	494
8	Extinction and Scattering Efficiencies for Composite Grains	497
	Figs. F1–F8: $Q_{ext}(\lambda^{-1})$, $Q_{sca}(\lambda^{-1})$ for composite grains with iron cores.	498
	Figs. G1–G16: $Q_{ext}(\lambda^{-1})$, $Q_{sca}(\lambda^{-1})$ for composite grains with graphite cores.	500
	Index	505

