

TABLE OF CONTENTS

Introduction	ix
I. Energy estimates for S^1-valued maps	
1. An auxiliary linear problem	1
2. Variants of Theorem I.1	6
3. S^1 -valued harmonic maps with prescribed isolated singularities. The canonical harmonic map	10
4. Shrinking holes. Renormalized energy	16
II. A lower bound for the energy of S^1-valued maps on perforated domains	31
III. Some basic estimates for u_ε	
1. Estimates when $G = B_R$ and $g(x) = x/ x $	42
2. An upper bound for $E_\varepsilon(u_\varepsilon)$	44
3. An upper bound for $\frac{1}{\varepsilon^2} \int_G (u_\varepsilon ^2 - 1)^2$	45
4. $ u_\varepsilon \geq 1/2$ on "good discs"	46
IV. Towards locating the singularities: bad discs and good discs	
1. A covering argument	48
2. Modifying the bad discs	49
V. An upper bound for the energy of u_ε away from the singularities	
1. A lower bound for the energy of u_ε near a_j	53
2. Proof of Theorem V.1	54

CONTENTS

VI. u_{ε_n} converges: u_* is born!	
1. Proof of Theorem VI.1	58
2. Further properties of u_* : singularities have degree one and they are not on the boundary	60
VII. u_* coincides with THE canonical harmonic map having singularities (a_j)	65
VIII. The configuration (a_j) minimizes the renormalized energy W	
1. The general case	76
2. The vanishing gradient property and its various forms	82
3. Construction of critical points of the renormalized energy	93
4. The case $G = B_1$ and $g(\theta) = e^{i\theta}$	95
5. The case $G = B_1$ and $g(\theta) = e^{di\theta}$ with $d \geq 2$	97
IX. Some additional properties of u_ε	
1. The zeroes of u_ε	100
2. The limit of $\{E_\varepsilon(u_\varepsilon) - \pi d \log \varepsilon \}$ as $\varepsilon \rightarrow 0$	101
3. $\int_G \nabla u_\varepsilon ^2$ remains bounded as $\varepsilon \rightarrow 0$	103
4. The bad discs revisited	104
X. Non minimizing solutions of the Ginzburg-Landau equation	
1. Preliminary estimates; bad discs and good discs	107
2. Splitting $ \nabla v_\varepsilon $	109
3. Study of the associated linear problems	112
4. The basic estimates: $\int_G \nabla v_\varepsilon ^2 \leq C \log \varepsilon $ and $\int_G \nabla v_\varepsilon ^p \leq C_p$ for $p < 2$	119
5. v_{ε_n} converges to v_*	125
6. Properties of v_*	132

CONTENTS

XI. Open problems	137
Appendix I. Summary of the basic convergence results in the case where $\deg(g, \partial G) = 0$	142
Appendix II. Radial solutions	145
Appendix III. Quantization effects for the equation $-\Delta v = v(1 - v ^2)$ in \mathbb{R}^2	147
Appendix IV. The energy of maps on perforated domains revisited	148
BIBLIOGRAPHY	154
INDEX	159