

Table of Contents

| | |
|--|-----|
| Preface | xii |
| Scientific Organising Committee | xiv |
| Introductory review: The Cosmic Distance Scale: Past, Present, and Future | |
| The cosmic distance scale and H_0 : Past, present, and future | 3 |
| <i>W. L. Freedman</i> | |
| Dedication in honour of Allan Sandage | |
| Allan Sandage and the distance scale | 13 |
| <i>G. A. Tammann & B. Reindl</i> | |
| Session I: Resolved stars in the Milky Way | |
| The distance to the Galactic Center | 29 |
| <i>S. Gillessen, F. Eisenhauer, T. K. Fritz, O. Pfuhl, T. Ott, & R. Genzel</i> | |
| The Gould's Belt Distances Survey | 36 |
| <i>L. Loinard</i> | |
| Adjusting the distance scales of old and intermediate-age tracers via statistical parallaxes | 44 |
| <i>A. K. Dambis</i> | |
| The PARSEC program: a large sample of brown dwarf trigonometric parallaxes | 48 |
| <i>A. H. Andrei, R. L. Smart, B. Bucciarelli, J. L. Penna, F. Marocco, M. G. Lattanzi, M. Crosta, & R. Teixeira</i> | |
| Direct distance determination using parallax: Techniques, promises and limitations | 52 |
| <i>L. Lindegren</i> | |
| Toward a VLBI resolution of the Pleiades distance controversy | 60 |
| <i>C. Melis, M. J. Reid, A. J. Mioduszewski, J. R. Stauffer, & G. C. Bower</i> | |
| A new assessment of the kinematic distance to the Pleiades: based on radial velocities and proper motions only | 66 |
| <i>S. Röser & E. Schilbach</i> | |
| Binaries and distances | 70 |
| <i>D. Pourbaix, F. Arenou, J.-L. Halbwachs, & C. Siopis</i> | |
| Determining distances to stars statistically from photometry | 74 |
| <i>H. J. Newberg</i> | |
| Astrometric observations of neutron stars | 82 |
| <i>S. Chatterjee</i> | |

| | |
|--|----|
| Spectroscopic distances to late-type stars | 83 |
| <i>M. Bergemann, A. Serenelli, & G. Ruchti</i> | |
| Parallaxes of metal-poor main-sequence stars | 87 |
| <i>B. Chaboyer, G. Feiden, G. F. Benedict, B. E. McArthur, T. E. Harrison, A. McWilliam, E. P. Nelan, R. J. Patterson, & A. Sarajedini</i> | |
| VLBI astrometry of the semi-regular variable star RX Bootis | 91 |
| <i>T. Kamezaki, A. Nakagawa, T. Omodaka, T. Handa, H. Imai, D. Tafoya, M. Matsui, Y. Nishida, T. Nagayama, M. Honma, H. Kobayashi, T. Miyaji, & M. Takeuti</i> | |
| The outer rotation curve project with VERA: Trigonometric parallax of IRAS 05168+3634 | 95 |
| <i>N. Sakai, M. Honma, H. Nakanishi, H. Sakanoue, T. Kurayama, & the VERA collaboration</i> | |

Session II: From the Milky Way to the Local Group

| | |
|---|-----|
| RR Lyrae stars: prime calibrators of the first rung of the distance ladder | 101 |
| <i>C. Cacciari</i> | |
| Cepheids and other variable stars and the distance to the Galactic Centre | 109 |
| <i>N. Matsunaga</i> | |
| Classical Cepheids, what else? | 116 |
| <i>G. Bono, L. Inno, N. Matsunaga, K. Genovali, B. Lemasle, F. Primas, & M. Romaniello</i> | |
| Physical parameters for RR Lyrae stars in the SDSS filter system | 126 |
| <i>M. Catelan, G. I. Torrealba, C. Cáceres, H. A. Smith, N. De Lee, & M. Fitzgerald</i> | |
| Distance estimates in the Milky Way using tidal streams | 130 |
| <i>D. S. M. Fantin</i> | |
| Wesenheit function for Galactic Cepheids: Application to the projection factors | 134 |
| <i>C.-C. Ngeow, H. Neilson, N. Nardetto, & M. Marengo</i> | |
| Cepheid distances from the Baade-Wesselink method | 138 |
| <i>W. Gieren, J. Storm, N. Nardetto, A. Gallenne, G. Pietrzynski, P. Fouqué, T. G. Barnes, & D. Majaess</i> | |
| Evolutionary properties of stellar standard candles: Red clump, AGB clump and white dwarfs | 145 |
| <i>M. Salaris</i> | |
| The Galactic Cepheid period-luminosity relation revisited using <i>bona fide</i> cluster Cepheids | 153 |
| <i>R. I. Anderson, N. Mowlavi, & L. Eyer</i> | |
| Circumstellar envelopes of Cepheids: a possible bias affecting the distance scale? | 157 |
| <i>P. Kervella, A. Gallenne, & A. Mérand</i> | |

| | |
|--|-----|
| The effects of physics and chemical-abundance uncertainties on the properties of lower-mass stars that are used as standard candles..... | 161 |
| <i>D. A. Vandenberg</i> | |
| A precise and accurate distance to the Large Magellanic Cloud from late-type eclipsing-binary systems..... | 169 |
| <i>G. Pietrzyński, W. Gieren, D. Graczyk, I. Thompson, B. Pilecki, N. Nardetto, R.-P. Kudritzki, F. Bresolin, G. Bono, P. Prada Moroni, P. Konorski, M. Gorski, J. Storm, R. Smolec, & P. Karczmarek</i> | |
| Eclipsing binary distances to the edge of the Local Group..... | 173 |
| <i>A. Z. Bonanos</i> | |
| The Araucaria Project: the Baade–Wesselink projection factor of pulsating stars and interferometric contribution to the eclipsing binaries technique of distance determination | 179 |
| <i>N. Nardetto, M. Chalouf, G. Guiglion, D. Mourard, W. Gieren, G. Pietrzyński, & D. Graczyk</i> | |
| Toward improving the accuracy of Cepheid distances through parallax of pulsation | 183 |
| <i>A. Mérand, P. Kervella, J. P. Aufdenberg, & A. Gallenne</i> | |
| The period–radius relation of classical Cepheids and the problem of their mode identification | 187 |
| <i>M. Sachkov</i> | |
| Galactic structure from trigonometric parallaxes of star-forming regions | 188 |
| <i>M. J. Reid</i> | |
| VERA status and results..... | 194 |
| <i>H. Kobayashi & the VERA collaboration</i> | |
| The Baade–Becker–Wesselink technique and the fundamental astrophysical parameters of Cepheids | 195 |
| <i>A. S. Rastorguev, A. K. Dambis, M. V. Zabolotskikh, L. N. Berdnikov, & N. A. Gorynya</i> | |
| From supergiant stars to galaxies: The path to extragalactic distances | 203 |
| <i>F. Bresolin & R.-P. Kudritzki</i> | |
| Asymptotic giant branch variables as extragalactic distance indicators | 209 |
| <i>P. A. Whitelock</i> | |
| The period–luminosity relation of Mira variable stars | 217 |
| <i>A. Nakagawa, T. Kurayama, T. Omodaka, T. Kamezaki, Y. Nishida, & M. Matsui</i> | |
| Maximum-likelihood implementation of the tip-of-the-red-giant-branch method . | 218 |
| <i>D. Makarov</i> | |
| The distance to the Small Magellanic Cloud from eclipsing binaries..... | 222 |
| <i>D. Graczyk, G. Pietrzyński, B. Pilecki, I. B. Thompson, W. Gieren, P. Konorski, A. Udalski, & I. Soszyński</i> | |

| | |
|---|-----|
| Rotational parallaxes of nearby galaxies | 226 |
| <i>A. Brunthaler</i> | |
| Application of the Direct Distance Estimation procedure to eclipsing binaries in star clusters..... | 227 |
| <i>E. F. Milone & S. J. Schiller</i> | |
| Distances, depth structure, and kinematics of the Magellanic Clouds from disentangling spectra of eclipsing binaries and Cepheids | 231 |
| <i>P. Hadrava, S. Štefl, R. Klement, & C. Martayan</i> | |
| The distance to M31 in the era of precision cosmology | 235 |
| <i>D. Valls-Gabaud</i> | |
| Multiparametric scaling relations for dwarf irregular galaxies in different environments..... | 236 |
| <i>M. E. Sharina, V. E. Karachentseva, & D. I. Makarov</i> | |
| The tip-of-the-red-giant-branch distance indicator and the structure of the nearest galaxy groups | 240 |
| <i>L. Makarova & D. Makarov</i> | |
| Session III: Reaching Virgo cluster distances and beyond | |
| The planetary nebula luminosity function | 247 |
| <i>R. Ciardullo</i> | |
| Measuring the Hubble constant with observations of water-vapor megamasers .. | 255 |
| <i>J. Braatz, M. Reid, C.-Y. Kuo, V. Impellizzeri, J. Condon, C. Henkel, K. Y. Lo, J. Greene, F. Gao, & W. Zhao</i> | |
| The Virgo Cluster | 262 |
| <i>J. Mould</i> | |
| The 6dFGS peculiar velocity field..... | 269 |
| <i>C. M. Springob, C. Magoulas, M. Colless, D. H. Jones, L. Campbell, J. Lucey, J. Mould, & P. Erdoğdu</i> | |
| The Carnegie Hubble Program: From parallaxes to the Tully–Fisher relation | 274 |
| <i>V. Scowcroft, W. L. Freedman, B. F. Madore, A. Monson, S. E. Persson, M. Seibert, C. Burns, & J. R. Rigby</i> | |
| Ultralong-period Cepheids: a possible primary distance indicator? | 282 |
| <i>G. Fiorentino, F. Annibali, G. Clementini, R. Contreras Ramos, M. Marconi, I. Musella, A. Saha, M. Tosi, A. Aloisi, & R. van der Marel</i> | |
| The planetary nebulae luminosity function and distances to the Virgo, Hydra I, and Coma Clusters..... | 287 |
| <i>M. Arnaboldi, A. Longobardi, O. Gerhard, & S. Okamura</i> | |
| Integrated photometric redshifts and SED fitting as tool for galaxy evolution studies | 292 |
| <i>R. Kotulla</i> | |
| On the scaling relations of disk galaxies..... | 296 |
| <i>R. Giovanelli</i> | |

| | |
|---|-----|
| Cosmic distances from surface brightness fluctuations | 304 |
| <i>J. P. Blakeslee</i> | |
| The 2MASS Tully–Fisher Survey: Mapping the mass in the Universe | 312 |
| <i>T. Hong, L. Staveley-Smith, K. Masters, C. Springob, L. Macri, B. Koribalski, H. Jones, & T. Jarrett</i> | |
| Session IV: Cosmological distances | |
| Measuring H_0 from the 6dF Galaxy Survey and future low-redshift surveys | 319 |
| <i>M. Colless, F. Beutler, & C. Blake</i> | |
| Visualization of structures and cosmic flows in the local Universe | 323 |
| <i>D. Pomarède, H. Courtois, & R. B. Tully</i> | |
| Type Ia supernovae as extragalactic distance indicators. | 327 |
| <i>B. P. Schmidt</i> | |
| Improving Type Ia supernova characterization through multiwavelength studies of Type Ia supernova host galaxies | 328 |
| <i>B. Tucker, B. P. Schmidt, & P. Garnavich</i> | |
| The heterogeneity of Type Ia supernova progenitor systems and their use as cosmic distance indicators | 329 |
| <i>P. Ricker, K.-C. Pan, & R. Taam</i> | |
| Testing the distance-duality relation to probe the morphology of galaxy clusters | 330 |
| <i>X.-L. Meng</i> | |
| Accurate cosmology from gravitational-lens time delays. | 331 |
| <i>S. H. Suyu</i> | |
| Measurement of the cosmological distance scale using X-ray and Sunyaev–Zel’dovich effect observations of galaxy clusters | 339 |
| <i>M. Bonamente, J. Carlstrom, E. Leitch, M. Joy, D. Marrone, A. Mantz, S. Muchevoj, T. Plagge, & E. Reese</i> | |
| Hubble constant, lensing, and time delay in TeVeS. | 344 |
| <i>Y. Tian, C.-M. Ko, & M.-C. Chiu</i> | |
| Summary | |
| Advancing the physics of cosmic distances: Conference summary | 351 |
| <i>R. de Grijs</i> | |
| Posters | |
| <i>Gaia</i> contribution to the low-redshift supernova population | 363 |
| <i>N. Blagorodnova, N. A. Walton, L. Wyrzykowski, & S. Hodgkin</i> | |
| Distance estimates to five open clusters based on 2MASS data of red clump giants | 367 |
| <i>L. Chen & X. Gao</i> | |
| Near-infrared surface brightness fluctuation measurements with the <i>Hubble Space Telescope</i> ’s WFC3/IR channel | 371 |
| <i>H. Cho, J. B. Jensen, J. P. Blakeslee, B. S. French, H.-c. Lee, & Y.-W. Lee</i> | |

| | |
|--|-----|
| Planetary nebulae and determination of the bulge–disk boundary | 375 |
| <i>R. D. D. Costa, O. Cavichia, & W. J. Maciel</i> | |
| Revisiting the absolute-magnitude calibration of F-type supergiants and bright giants as a function of the equivalent width of the OI λ 7774Å triplet | 379 |
| <i>A. K. Dambis</i> | |
| Recalibration of the <i>uvby</i> β photometry-based distance scale of OB-type stars using the revised <i>Hipparcos</i> parallaxes | 382 |
| <i>A. K. Dambis</i> | |
| Trigonometric parallaxes in the TW Hydrae Association | 386 |
| <i>C. Ducourant, R. Teixeira, G. Chauvin, A. Krone-Martins, J. F. Le Campion, P. Galli, I. Song, & B. Zuckerman</i> | |
| Kinematic distances of pre-main-sequence stars in the Lupus star-forming region | 390 |
| <i>P. A. B. Galli, R. Teixeira, C. Ducourant, & C. Bertout</i> | |
| Global survey of star clusters in the Milky Way: The open cluster distance scale | 394 |
| <i>N. V. Kharchenko, A. E. Piskunov, S. Röser, E. Schilbach, & R.-D. Scholz</i> | |
| Determination of the fundamental properties of an M31 globular cluster from main- sequence photometry | 398 |
| <i>J. Ma</i> | |
| Maximum-likelihood fitting of the 6dFGS peculiar velocities | 402 |
| <i>C. Magoulas, C. Springob, M. Colless, D. H. Jones, L. Campbell, J. Lucey, & J. Mould</i> | |
| Statistical analysis of the determinations of the Sun’s Galactocentric distance . | 406 |
| <i>Z. Malkin</i> | |
| 3D motions of 6.7 GHz methanol masers and effects of the Galactic bar | 410 |
| <i>N. Matsumoto, M. Honma, & the VERA collaboration</i> | |
| Improving distance estimates to nearby bright stars: Combining astrometric data from <i>Hipparcos</i> , <i>Nano-JASMINE</i> and <i>Gaia</i> | 414 |
| <i>D. Michalik, L. Lindegren, D. Hobbs, U. Lammers, & Y. Yamada</i> | |
| VERA observations of the Galactic star-forming regions ON1 and ON2N | 418 |
| <i>T. Nagayama & VERA project members</i> | |
| The kinematics and chemistry of the Sagittarius streams | 422 |
| <i>W. Shi, Y. Chen, K. Carrell, & G. Zhao</i> | |
| Using L dwarfs with distances to test models | 425 |
| <i>Y. Wang, H. R. A. Jones, R. L. Smart, & Z. Shao</i> | |
| Scientific goals of <i>Nano-JASMINE</i> | 429 |
| <i>Y. Yamada, S. Fujita, N. Gouda, Y. Kobayashi, T. Hara, R. Nishi, S. Yoshioka, S. Hozumi, & the JASMINE Working Group</i> | |
| The scientific goal of the Japanese small astrometric satellite, <i>Small-JASMINE</i> . | 433 |
| <i>T. Yano, N. Gouda, Y. Kobayashi, T. Tsujimoto, Y. Niwa, & Y. Yamada</i> | |

| | |
|--|-----|
| Evaluation of microlens distributions in gravitationally lensed systems based on accurate radio observations | 437 |
| <i>A. F. Zakharov, S. Simić, L. Č. Popović, & P. Jovanović</i> | |
| A preliminary systematic search for red-clump stars in Galactic open clusters based on 2MASS data | 441 |
| <i>X. Zhang, L. Chen, & Z. Li</i> | |
| Galactocentric distance of the Sun derived from kinematic data | 444 |
| <i>Z. Zhu & M. Shen</i> | |
| Author index | 449 |
| Object index | 453 |
| Subject index | 459 |