

Contents

1	Historical perspective	1
2	Electronic states, phonons, and electron-phonon interaction	5
2.1	Adiabatic approximation: Hamiltonian	5
2.2	Adiabatic approximation and non-adiabaticity: Born–Oppenheimer and “crude” approaches	6
2.3	Electron–phonon coupling	9
2.4	Electron–phonon interaction and renormalization of normal parameters	11
2.5	The “Migdal” theorem	15
2.6	Polaronic states	16
2.6.1	Concept	16
2.6.2	Dynamic polaron	17
3	Phonon mechanism	20
3.1	Superconductivity as a “giant” non-adiabatic phenomenon	20
3.2	The BCS model	21
3.3	Phonon mechanism: main equations	22
3.4	Critical temperature	26
3.4.1	Weak coupling	26
3.4.2	Intermediate coupling ($\lambda \lesssim 1.5$)	28
3.4.3	Coulomb interaction	29
3.4.4	Very strong coupling	31
3.4.5	The general case	33
3.4.6	About an upper limit of T_c	35
3.5	Properties of superconductors with strong coupling	36
3.6	The Van Hove scenario	39
3.7	Bipolarons: BEC versus BCS	39
3.8	Superconducting semiconductors	40
3.9	Polaronic effect and its impact on T_c	42
3.9.1	Double-well structure	42
3.9.2	Superconducting state	44
4	Electronic mechanisms	47
4.1	The Little model	47
4.2	“Sandwich” excitonic mechanism	50
4.3	Three-dimensional systems: electronic mechanism	50
4.4	Plasmons	52
4.4.1	Plasmons in layered systems: dispersion law and “electronic sound”	53

4.4.2	Plasmons in layered conductors: pairing	57		
4.4.3	The 3D case: “demons”	58		
5	Magnetic mechanism	59		
5.1	Introduction	59		
5.1.1	Localized versus itinerant aspects of the cuprates	60		
5.2	Fermi liquid-based theories	62		
5.2.1	The spin-bag model of Schrieffer, Wen, and Zhang (1989)	62		
5.2.2	The t - J model (Emery, 1987; Zhang and Rice, 1988)	66		
5.2.3	Two-dimensional Hubbard model studies by Monte Carlo techniques	70		
5.2.4	Spiral phase of a doped quantum antiferromagnet (Shraiman and Siggia, 1988–89)	77		
5.2.5	Slave bosons	82		
5.3	Non-Fermi-liquid models	85		
5.3.1	The resonant valence bond (RVB) model and its evolution	85		
5.3.2	Anyon models and fractional statistics	86		
5.4	Conclusions	87		
6	Experimental methods: Spectroscopic	88		
6.1	Tunneling spectroscopy	88		
6.1.1	Experimental method	88		
6.1.2	Energy gap and transition temperature	90		
6.1.3	Inversion of the gap equation and $\alpha^2 F(\Omega)$	91		
6.1.4	Electron-phonon coupling parameter λ	94		
6.2	Scanning tunneling microscopy and spectroscopy	96		
6.3	Infrared spectroscopy	97		
6.4	Ultrasonic attenuation	99		
6.5	Angle-resolved photoemission	100		
6.6	Muon spin resonance (μ SR)	100		
6.6.1	μ SR studies of superconductivity	102		
7	Multigap superconductivity	103		
7.1	Multigap superconductivity: general picture	103		
7.2	Critical temperature	104		
7.3	Energy spectrum	105		
7.4	Properties of two-gap superconductors	108		
7.4.1	Penetration depth; surface resistance	108		
7.4.2	Strong magnetic field: Ginzburg–Landau equations for a multigap superconductor	110		
7.4.3	Heat capacity	111		
7.4.4	Experimental data	111		
7.5	Induced two-band superconductivity	112		
7.6	Symmetry of the order parameter and multiband superconductor	113		
8	Induced superconductivity: proximity effect	114		
8.1	Proximity “sandwich”	114		
8.2	Critical temperature	115		
8.3	Proximity effect versus the two-gap model	119		
8.4	Pair-breaking: gapless superconductivity	119		
9	Isotope effect	122		
9.1	General remarks	122		
9.2	Coulomb pseudopotential	122		
9.3	Multi-component lattice	123		
9.4	Anharmonicity	123		
9.5	Isotope effect in proximity systems	124		
9.6	Magnetic impurities and isotope effect	125		
9.7	Polaronic effect and isotope substitution	126		
9.8	Penetration depth: isotopic dependence	128		
10	Cuprate superconductors	131		
10.1	History	131		
10.2	Structure of the cuprates	132		
10.3	Preparation of bulk and film cuprates	133		
10.4	Properties of the cuprates	134		
10.4.1	Phase diagram	134		
10.4.2	Critical field H_{c2}	135		
10.4.3	Two-gap spectrum	136		
10.4.4	Symmetry of the order parameter	136		
10.5	Isotope effect	138		
10.5.1	Polaronic state	138		
10.5.2	Isotopic dependence of the penetration depth	140		
10.6	Mechanism of high T_c	140		
10.7	Proposed experiment	145		
11	Inhomogeneous superconductivity and the “pseudogap” state of novel superconductors	147		
11.1	“Pseudogap” state: main properties	148		
11.1.1	Anomalous diamagnetism above T_c	148		
11.1.2	Energy gap	150		
11.1.3	Isotope effect	152		
11.1.4	“Giant” Josephson effect	152		
11.1.5	Transport properties	153		
11.2	Inhomogeneous state	154		
11.2.1	Qualitative picture	154		
11.2.2	The origin of inhomogeneity	155		
11.2.3	Percolative transition	156		
11.2.4	Inhomogeneity: experimental data	156		
11.3	Energy scales	157		
11.3.1	Highest-energy scale (T^*)	158		

11.3.2 Diamagnetic transition (T_c^*)	158	13.4 Granular superconductors	202
11.3.3 Resistive transition (T_c)	159	13.5 Sr_2RuO_4 : a very novel superconductor	203
11.4 Theory	159	13.6 Ruthenium cuprates	204
11.4.1 General equations	160	13.7 Intercalated nitrides: self-supported superconductivity	205
11.4.2 Diamagnetism	160		
11.4.3 Transport properties; “giant” Josephson effect	162		
11.4.4 Isotope effect	166		
11.5 Other systems	167	14 Organic superconductivity	206
11.5.1 Borocarbides	167	14.1 History	206
11.5.2 Granular superconductors; Pb+Ag system	167	14.2 Organic superconductors: structure, properties	207
11.6 Ordering of dopants and potential for room-temperature	168	14.3 Intercalated materials	210
superconductivity	168	14.4 Fullerides	212
11.7 Remarks	171	14.5 Small-scale organic superconductivity	213
		14.6 Pair correlation in aromatic molecules	214
12 Manganites	172	15 Pairing in nanoclusters: nano-based superconducting	218
12.1 Introduction	172	tunneling networks	218
12.2 Electronic structure and doping	173	15.1 Clusters: shell structure	218
12.2.1 Structure	173	15.2 Pair correlation	220
12.2.2 Magnetic order	176	15.2.1 Qualitative picture	220
12.2.3 Double-exchange mechanism	176	15.2.2 Main equations: critical temperature	222
12.2.4 Colossal magnetoresistance (CMR)	177	15.2.3 Energy spectrum; fluctuations	225
12.3 Percolation phenomena	178	15.3 How to observe the phenomenon?	226
12.3.1 Low doping: transition to the ferromagnetic state	178	15.4 Cluster-based tunneling network: macroscopic superconductivity	227
at low temperatures		15.5 Cluster crystals	228
12.3.2 Percolation threshold	179		
12.3.3 Increase in temperature and percolative transition	180	Appendices	229
12.3.4 Experimental data	181	Appendix A: Diabatic representation	229
12.3.5 Large doping	182	Appendix B: Dynamic Jahn-Teller effect	231
12.4 Main interactions: Hamiltonian	183	References	233
12.5 Ferromagnetic metallic state	184	Index	255
12.5.1 Two-band spectrum	184		
12.5.2 Heat capacity	186		
12.5.3 Isotope substitution	187		
12.5.4 Optical properties	189		
12.6 Insulating phase	190		
12.6.1 Parent compound	190		
12.6.2 Low doping: polarons	191		
12.7 Metallic A-phase: $S-N-S$ Josephson effect	193		
12.7.1 Magnetic structure	193		
12.7.2 Josephson contact with the A-phase barrier	193		
12.8 Discussion: manganites versus cuprates	195		
13 Novel superconducting systems	197		
13.1 Fe-based pnictide and chalcogenide superconductors	197		
13.2 Magnesium diboride: MgB_2	199		
13.3 A-15 structure superconductors	201		