Contents

| On the definition of a positive frequency part for a quantized scalar field in | | | | | | | |
|--|-----|--|--|--|--|--|--|
| several Robertson-Walker universes H. Nariai aad T. Azuma | 1 | | | | | | |
| A gauge invariant approach to two-dimensional quantum chromodynamics | | | | | | | |
| K. Kikkawa | 15 | | | | | | |
| A unified potential for mesons and baryons | | | | | | | |
| R. K. Bhaduri, L. E. Cohler and Y. Nogami | 30 | | | | | | |
| Longitudinal polarization of ¹² B in muon capture reaction | | | | | | | |
| H. Ami, M. Kobayashi, H. Ohtsubo and M. Morita | 45 | | | | | | |
| Radiative pion capture on ¹² C N. Ohtsuka and H. Ohtsubo | 62 | | | | | | |
| Production of charged pions by high energy photon on nuclei | | | | | | | |
| Y. Futami and J. Suzumura | 78 | | | | | | |
| Short range part of baryon-baryon interaction in a quark model. I | | | | | | | |
| ——Formulation—— M. Oka and K. Yazaki | 91 | | | | | | |
| Inelastic scattering of pions on ¹² C | | | | | | | |
| F. M. Toyama, Y. Hatsuda and Y. Sakamoto | 107 | | | | | | |
| On the bound state of the trineutron by solving the Faddeev equation in | | | | | | | |
| coordinate space Y. Sunami | 116 | | | | | | |
| Microscopic approach to two-nucleon absorption of in-flight pions in nuclei | | | | | | | |
| T. Takaki | 124 | | | | | | |
| Relativistic corrections to perturbation theory for pionic and kaonic atoms | | | | | | | |
| M. Leon and R. Seki | 137 | | | | | | |
| Nuclear effective interactions and spin excitations T. Kohmura | 145 | | | | | | |
| Damping state method and kaonic bound states M. Atarashi | 163 | | | | | | |
| A separable potential and off-shell scattering amplitude for an α - Λ | | | | | | | |
| interaction S. Oryu and R. Kircher | 176 | | | | | | |
| Continuous radiation spectra of positron annihilation in flight | | | | | | | |
| K. Shizuma, S. Hamanaka and Y. Yoshizawa | 180 | | | | | | |
| Ionization of atomic hydrogen by electron impact in the eikonal | | | | | | | |

| approximation with exchange T. Sekimura | 185 | | | | | |
|---|-----|--|--|--|--|--|
| Feasibility of using a high-Rydberg atom in a collision experiment as a | | | | | | |
| probe to scattering of an extremely slow electron by a rare-gas atom | | | | | | |
| K. Sasano and M. Matsuzawa | 197 | | | | | |
| Low-energy phaseshift formula for e±-atom elastic scattering | | | | | | |
| T. Sasakawa and J. Horáček | 203 | | | | | |
| Triple differential cross sections for the ionization of atomic hydrogen by | | | | | | |
| electron impact in the Coulomb-projected-eikonal approximation | | | | | | |
| WJ. Chen | 209 | | | | | |
| Eikonal approach to break-up processes of hydrogen-like ions by electron | | | | | | |
| impact: Exchange of electrons H. Kotegawa | 218 | | | | | |
| Spectral line shifts produced by electron collisions in HeII plasma | | | | | | |
| K. Yamamoto | 227 | | | | | |
| Interaction of ion-acoustic solitons in two-dimensional space | | | | | | |
| F. Kako and N. Yajima | 238 | | | | | |
| Physics of wave-particle interactions K. Nishikawa | 247 | | | | | |
| An energy integral and variational principle for plasma kinetic modes | | | | | | |
| SI. Itoh, K. Itoh and Y. Terashima | 267 | | | | | |
| Stiff-chain statistics in relation to the Brownian process A. Miyake | 272 | | | | | |
| Additional facilities of a conventional DBMS to support interactive | | | | | | |
| statistical analysis H. Ikeda and Y. Kobayashi | 279 | | | | | |
| Conjugate shake-up process of neon due to K-shell photoionization | | | | | | |
| T. Ishihara, J. Mizuno and T. Watanabe | 291 | | | | | |
| Scattering by a Coulomb plus spin-orbit interaction in the Glauber | | | | | | |
| approximation V. Franco | 297 | | | | | |
| Electron capture into partially stripped projectile ions | | | | | | |
| J. Eichler, A. Tsuji and T. Ishihara | 302 | | | | | |
| Energy loss by slow electron and by slow atoms in a molecular gas | | | | | | |
| I. Shimamura | 310 | | | | | |
| Delta-hole formalism of pion-nucleus single charge exchange scattering | | | | | | |
| M. Hirata | 314 | | | | | |
| p-state contribution to the Λ well depth | | | | | | |
| M. Kimura, Y. Takahashi and E. Satoh | 321 | | | | | |
| | | | | | | |

xii